



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Multiplying these sides respectively by 16, 9 and 4, we obtain the three triangles whose sides are respectively 41, 9, 40; 34, 16, 36, and 29, 21, 20.

If we desire to determine a particular kind of triangle, having for instance, the difference between their base and perpendiculars = 1; put $a = \frac{1}{5}$, and multiply the resulting sides by 25, and we get 120, 119, 169. Put $a = \frac{4}{17}$, and multiply by $\frac{289}{2}$, and we get 697, 696, 985. Put $a = \frac{159}{70}$, and multiply by 4900, and we get 23660, 23661, 33461, &c.

This last class of triangles I consider among the most interesting.

THE CORRECT METHOD OF LEAST SQUARES.—BY R. J. ADCOCK, MONMOUTH, ILL.—1. The point determined by two given points is their center of gravity. The point determined by any number of given points is their center of gravity. For combining them in sets of two, so as to take each point twice, gives the same number of other points, which being combined continually, as before, finally terminates at the center of gravity of the given points.

2. Hence a straight line which is to pass through a given point, having its direction to be determined by other given points, must pass through their center of gravity; because that is the point determined by the other given points.

3. Hence also, a straight line determined by any given points, passes through their center of gravity, and has such a direction that it passes through the two most remote centers of gravity of the two sets of points on opposite sides of the common center of gravity; and is that principal axis of least moment of inertia of the given points.

4. Hence in general, any point, line or surface, determined by any given points, is that point, line or surface which has the sum of the squares of the normals to it from the given points, a minimum. And is the point, line or surface which has the mean position of all that can be taken.

The common application of the method is incorrect, not agreeing with 2, 3, and 4.

ANSWER TO PROF. BROOKS' QUERY.—No rigorous demonstration can be given of the formula mentioned, it must be regarded as empirical.

The function $\phi(x)$, which denotes the number of primes less than x , is discontinuous, and the problem of finding functions representing it approximately is an indeterminate one. Legendre was the first to give a formula of this kind, (*Essai sur la Theorie des Nombres*, 2d Edition, Paris, 1808, p. 394, or 3d Edition, Vol. II, p. 65); it is

$$N = \frac{x}{\log x - 1.08366},$$

$\log x$ being Napierian. The constant 1.08366 evidently has been assumed so as to represent as well as possible the actual count of primes below